

Motivational Interviewing With Parents for Obesity: An RCT

Amir H. Pakpour, PhD^a, Paul Gellert, PhD^b, Stephan U. Dombrowski, PhD^c, Bengt Fridlund, PhD^d

abstract

BACKGROUND AND OBJECTIVE: Motivational interviewing (MI) has been shown to be an effective strategy for targeting obesity in adolescents, and parental involvement is associated with increased effectiveness. The aim of this study was to evaluate and compare the role of parental involvement in MI interventions for obese adolescents.

METHODS: A total of 357 Iranian adolescents (aged 14–18 years) were randomized to receive an MI intervention or an MI intervention with parental involvement (MI + PI) or assessments only (passive control). Data regarding anthropometric, biochemical, psychosocial, and behavioral measures were collected at baseline and 12 months later. A series of intention-to-treat, 2-way repeated-measures analysis of covariance were performed to examine group differences in change in outcomes measures over the 12-month follow-up period.

RESULTS: Results revealed significant effects on most of the outcome parameters for MI + PI (eg, mean \pm SD BMI z score: 2.58 ± 0.61) compared with the passive control group (2.76 ± 0.70 ; post hoc test, $P = .02$), as well as an additional superiority of MI + PI compared with MI only (2.81 ± 0.76 ; post hoc test, $P = .05$). This pattern was also shown for most of the anthropometric, biochemical, psychometric, and behavioral outcome variables.

CONCLUSIONS: MI with parental involvement is an effective strategy in changing obesity-related outcomes and has additional effects beyond MI with adolescents only. These findings might be important when administering MI interventions in school settings.



WHAT'S KNOWN ON THIS SUBJECT: Motivational interviewing (MI) has been found to increase the effectiveness of weight loss programs in obese children and adolescents. Although parental involvement seems to be linked to its effectiveness, strong conclusions cannot be drawn.

WHAT THIS STUDY ADDS: The present study found that MI with parental involvement is an effective strategy in changing obesity-related outcomes and has additional effects beyond MI with adolescents only. These findings are important when administering MI interventions in school settings.

^aSocial Determinants of Health Research Centre, Qazvin University of Medical Sciences, Qazvin, Iran;

^bCharité–Universitätsmedizin, Berlin, Germany; ^cUniversity of Stirling, United Kingdom; and ^dJönköping University, Jönköping, Sweden, Sweden

Dr Pakpour conceptualized and designed the study, conducted the initial analyses, and drafted the initial manuscript; Dr Gellert analyzed and interpreted the data, drafted the initial manuscript, and reviewed and revised the manuscript critically; Drs Dombrowski and Fridlund interpreted the data and reviewed and revised the manuscript critically; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

This trial has been registered at www.clinicaltrials.gov (NCT0218080).

www.pediatrics.org/cgi/doi/10.1542/peds.2014-1987

DOI: 10.1542/peds.2014-1987

Accepted for publication Dec 15, 2014

Address correspondence to Amir H. Pakpour, PhD, Social Determinants of Health Research Center, Qazvin University of Medical Sciences, Shahid Bahonar BLV, Qazvin, 3419759811, Iran.
E-mails: pakpour_amir@yahoo.com, apakpour@qums.ac.ir

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

Childhood obesity is a global health concern, and many obese adults were obese during childhood.^{1,2} Almost 10% of school-aged children worldwide are obese, and >60% of European children are overweight.^{3,4} In developing countries, childhood obesity is also reaching critical levels.⁵ In Iran, obesity among children and adolescents ranges from 7.1% to 12%.^{6,7} Childhood obesity increases complications such as skeletal/joint problems, social and mental health issues due to stigmatization, and reduced self-esteem.^{8,9} Prediabetes, hyperglycemia, and sleep apnea are increasingly prevalent among obese children.^{8,10} The prevention and treatment of obesity in children are priorities for public health worldwide.

Behavioral interventions targeting changes in physical activity and diet are effective for weight loss.^{11,12} Nearly 80% of pediatricians report frustration with modifying children's obesity, however.¹³ Patient motivation is associated with program adherence and predicts weight reduction.¹⁴ Including brief motivational techniques such as motivational interviewing (MI) in weight control interventions increases effectiveness.^{15,16}

MI is a client-centered clinical method to facilitate behavioral change.¹⁷ Originally developed as an intervention for addictive behaviors, MI is applied to a multitude of behavior change contexts.¹⁸ As part of MI, the participants' readiness to change is assessed and guides the discussions. The core principles of MI are: expressing empathy, developing a discrepancy, rolling with resistance, and supporting self-efficacy.^{18,19} MI invites participants to indicate their feelings, thoughts, and ambivalence about change, and it helps reaching decisional balance.^{17,20} Compared with other approaches such as education and nonspecific counseling, MI has been found to improve health behaviors and outcomes.¹⁹

Parental involvement in MI interventions²¹ can improve the effectiveness of school-based health programs. In a recent meta-analysis of MI interventions for health behavior change in adolescents, the overall effect size (Hedges' *g*) compared with other active treatment or no treatment was 0.28 (37 studies). Obesity interventions had a smaller effect size (*g* = 0.15; 12 studies). Regarding the intervention target/recipient,²² MI studies targeting both the adolescent and parent yielded the largest effect sizes (*g* = 0.59) and were more effective than conducting MI with either parent (*g* = 0.27) or child (*g* = 0.26) individually.²³

Although parental involvement is linked to effectiveness, conclusions about the appropriate intensity cannot be drawn due to the small number of studies and inconsistent results.^{24,25} Parenting skills (generic and specific to lifestyle behavior) were frequently present in effective weight control interventions,²⁴ and indirect methods to engage parents were commonly used, although direct approaches were more likely to result in positive outcomes.^{24,26}

Parental involvement requires further investigation with the use of randomized controlled trial (RCT) designs.²³ Although meta-analytical differences in effect sizes favor parental involvement, no existing RCT studies compare >2 of the 3 target groups (eg, MI, MI + PI, control) in the same trial experimentally.^{23,27–29}

The goal of the present RCT was to evaluate the effectiveness of an MI weight loss intervention for obese adolescents including parental involvement (MI + PI) versus MI without parental involvement (MI only) and a passive control group over a period of 12 months. We hypothesized that MI + PI would be effective in changing BMI and other relevant outcomes compared with a control condition. We further hypothesized that MI + PI would be

superior in changing weight loss outcomes compared with MI only.

METHODS

Participants and Recruitment

Obese adolescents (ie, BMI ≥95th percentile for age and gender³⁰) between 13 and 18 years of age were recruited from a local outpatient pediatric clinic in Qazvin, Iran. Exclusion criteria were taking weight-related medication, having a diagnosis of an eating disorder, being pregnant, and having clinical mental health conditions or psychosis. Eligible adolescents lived with a parent or adult caregiver who was prepared to be involved in treatment.

Study Design

The present study included a prospective RCT design in which patients were randomly assigned to a control group, MI with parental involvement, or MI without parental involvement (Fig 1). Anthropometric, biochemical, psychosocial, and behavioral measures were obtained at baseline and 12 months later.

All participants provided written informed consent; parental assent was also requested for adolescents aged <16 years. The ethics committee of the Qazvin University of Medical Sciences approved the study. A total of 409 adolescents were recruited between February 2011 and April 2013.

Power Analysis and Randomization Procedure

The sample size was calculated based on previous studies considering an effect size of 0.47, a power of 95%, and an α level of .05.³¹ Assuming a dropout rate of 15%, a sample size of 136 participants per condition was estimated.

To ensure adequate concealment of allocation, a research coordinator performed the randomization procedure by using a computer-generated




FIGURE 1
Flow chart of participants according to the Consolidated Standards of Reporting Trials.

randomization schedule/sequence (SPSS version 17.0, IBM SPSS Statistics, IBM Corporation, Armonk, NY). After recruitment and consenting procedures were completed, baseline data were collected, and a research coordinator assigned adolescents randomly to the 3 groups: control ($n = 119$), MI ($n = 119$), and MI + PI ($n = 119$). Adolescents could not be blinded to intervention allocation. Anthropometric measures at baseline and 12 months after randomization were taken by 2 assessors blinded to group allocation.

Intervention Groups

The MI intervention component targeted improved eating and physical activity behavior. The adolescents were encouraged to eat a variety of foods from each of the 4 major food groups and low-fat alternatives.³² They were also encouraged to achieve at least

60 minutes of moderate to vigorous intensity physical activity daily as recommended by the World Health Organization³³ (Supplemental Appendix).

MI Group With Parental Involvement

All sessions were delivered by 2 trained interventionists at the same time; each interventionist had different expertise (ie, registered dietitian and exercise specialist). All participants in the interventional groups attended pediatric clinics and received the weekly sessions lasting for 40 minutes. Both intervention groups received identical MI sessions (Supplemental Appendix). However, parents or guardians of participants from the MI + PI group ($n = 119$) received 1 additional session with >60 minutes in the clinic delivered by the same 2 interventionists at the end of the 6 sessions. An identical MI style was used for parents, focusing on the adolescents' weight, parents'

attitudes and behaviors regarding children's physical activity and dietary habits, parent monitoring, and supervision; the goal was to promote progress toward the child's intervention goals and attitudes by the parents.

MI Integrity

All MI sessions were audiotaped and quality checked by using the Motivational Interviewing Treatment Integrity (MITI) instrument.³⁴ Twenty-five percent of the interviews were selected randomly and assessed by 2 trained evaluators. The MITI has 2 components: global scores and behavior counts. Five global dimensions are rated on a 5-point Likert scale ranging from 1 (low) to 5 (high): evocation, collaboration, autonomy/support, direction, and empathy. A behavior count requires the coder to tally instances of particular interviewer behaviors, including providing information, MI-adherent behavior, MI-nonadherent behavior, simple reflections, complex reflections, and open and closed questions. In addition to the aforementioned scores, summary scores were computed as the following; global spirit rating (evocation + collaboration + autonomy/support)/3, percent complex reflections (number of complex reflections/number of total reflections) $\times 100$, percent open questions (number of open questions/number of total questions) $\times 100$, reflection-to-question ratio (number of total reflections/number of total questions) $\times 100$, and percent MI-adherent (number of MI-adherent statements/number of MI-adherent and MI-nonadherent statements) $\times 100$. As recommended according to the MITI, summary scores have applications for therapy coding. For ease of interpretation, we report summary scores for determining competence in MI.

The mean \pm SD scores were as follows: percent MI-adherent, 91.51 ± 16.12 ; percent open questions, 58.61 ± 13.20 ;

percent complex reflections, 49.37 ± 15.42 ; reflection-to-question ratio, 1.12 ± 0.38 ; and global spirit rating, 3.82 ± 1.03 . All scores were above proficiency according to the MITI recommendation, with the exception of the percent complex reflections, which was slightly below proficiency.³⁴

Measures

Primary outcomes were changes in BMI, the Child Dietary Self-Efficacy Scale (CDSS), the Weight Efficacy Lifestyle questionnaire (WEL), the Physical Exercise Self-Efficacy Scale (PES), and self-reported physical activity and diet. Secondary outcomes were changes in anthropometric measures, cholesterol, triglycerides, and body fat. All outcomes were assessed by 2 blinded and trained physicians. The blinded assessors passed a series of training courses for testing and interviewing according to the standard program protocol. Interrater reliability and interrater reliability were checked for both assessors before conducting the study.

Child Dietary Self-Efficacy Scale

The CDSS is a 15-item tool assessing dietary self-efficacy among children between 7 and 10 years of age. The measure begins with stems such as "How sure are you ..."³⁵ Responses are gathered on a 3-point Likert scale, ranging from "not sure" to "very sure." Responses ranged from -15 to 15, with higher scores indicating higher dietary self-efficacy. The CDSS had high internal consistency ($\alpha = .90$).

Weight Efficacy Lifestyle Questionnaire

The WEL is a 20-item tool assessing adolescents' confidence in their ability to lose weight. Items are scored on a 10-point Likert scale, ranging from 0 (not confident) to 9 (very confident), and items scores were averaged. The WEL is valid and reliable in weight management research.³⁶ The WEL had high internal consistency ($\alpha = .88$).

Psychometric properties of the Persian version of the WEL were assessed in Iranian adolescents, and the WEL was found to be a highly valid and reliable instrument.³⁷

Physical Exercise Self-Efficacy Scale

Adolescents' confidence in their ability to perform physical activity was measured by using the PES, a 5-item tool with responses scored on a 4-point Likert scale ranging from 0 (uncertain) to 9 (very certain). Cronbach's α for the PES was 0.92.

Diet

Diet was measured by using the Youth Adolescent Food Frequency Questionnaire (YAQ). The YAQ assesses diet among children and adolescents (aged 9–18 years) and is based on intake over the past year; it is valid and reliable against the 24-hour dietary recall adolescent populations.³⁸ The YAQ has 152 items and takes ~20 to 30 minutes to complete. The types and amounts of food consumed during the past 12 months were reported, and total energy intake (in kilocalories), total dietary fat, saturated fat, and servings per day of fruits, vegetables, snacks and desserts, milk, nondiet soda, and fried foods were analyzed. The Persian version of the YAQ was found to be valid and reliable for use in Iranian adolescents (A.H.P., Saffari M, Chen H. Psychometric evaluation of the Persian version of the Youth/Adolescent Questionnaire [YAQ] in Iranian obese adolescents, submitted manuscript).

Physical Activity

Physical activity was assessed by using a 7-day physical activity recall interview. This semi-structured interview is used to estimate an adolescent's time spent in physical activity, his or her strength, and flexibility activities for the 7 days before the interview. The total score for the physical activity recall interview was computed by using hours spent in physical activities of

moderate, hard, and very hard intensity. To estimate participants' energy expenditure (in kilocalories), the time spent in each activity for the past 7 days are multiplied by an intensity factor (1.5 for light intensity, 4 for moderate intensity, 6 for hard intensity, and 10 for very hard intensity).³⁹ The validity and reliability of the Persian version of the questionnaire have been well documented.⁴⁰

Objective Physical Activity

Objective physical activity was measured by using a GT3× monitor (ActiGraph LLC, Pensacola, FL), which is a 1-dimensional piezo-electric accelerometer. The GT3× is a light, small device that is a valid and reliable tool to assess physical activity.⁴¹ Due to financial restrictions, only 25 randomly selected adolescents in each group ($n = 75$) provided accelerometer data. Accelerometers were worn on the right hip for 7 days continuously except during showering or swimming. Adolescents who wore the device <8 hours per day or 4 days per week were excluded from the analysis. Sixty adolescents (~20 adolescents in each group) wore the device as instructed.

Pediatric Quality of Life Inventory

Quality of life was assessed by using the Pediatric Quality of Life Inventory 4.0 Short Form 15 (PedsQL 4.0 SF15). The PedsQL 4.0 SF15 has 15 items covering 4 dimensions: the physical functioning scale (5 items), the emotional functioning scale (4 items), the social functioning scale (3 items), and the school functioning scale (3 items). The physical health summary score is the same as the physical functioning scale, whereas the psychosocial health summary score is computed by the mean items of the emotional, social, and school functioning subscales. Items are reverse scored and linearly transformed to a scale of 0 to 100, with higher scores indicating better

quality of life. A recent study supports the feasibility, reliability, and validity of the Iranian version of the PedsQL 4.0 SF15 among Iranian children and adolescents.⁴²

Anthropometric, Body Composition, and Biologic Measurements

All clinical measures were performed at baseline and at a 12-month follow-up. Serum triglyceride and total cholesterol levels were assessed by using an autoanalyzer (Hitachi 704, Boehringer Mannheim, Mannheim, Germany).

Height for both adolescents and parents was measured to the nearest 0.1 cm after removing shoes. Height was measured by using a stadiometer (Seca Model 207, Seca, Hamburg, Germany). Weight for adolescents and parents was also measured to the nearest 0.1 kg on calibrated digital scales. BMI was calculated by dividing body weight by squared height in meters. As recommended by the World Health Organization, the growth reference chart based on z scores provides useful information on young people's nutritional status and growth. A BMI z score or SD score indicates how many units (of the SD) a child's BMI is above or below the average BMI value for his or her age group and gender.⁴³ The BMI percentile was calculated according to the Centers for Disease Control and Prevention's age- and gender-specific reference norms.³⁰

Waist circumference was measured midway between the lowest rib and the superior border of the iliac crest with an inelastic measuring tape at the end of normal expiration to the nearest 0.1 cm.

Percentage of body fat was measured by using the dual-energy x-ray absorptiometry (DXA). The DXA is a valid measure that provides a more accurate assessment of body composition than body weight alone. To further assess the adiposity of the adolescents, a bioelectrical impedance analysis (BIA) was

conducted. The BIA has been widely used to estimate body composition with simple, quick, and noninvasive technology.⁴⁴ Despite a considerable correlation between DXA and BIA in adult studies,^{45,46} results of pediatric studies have shown that these methods cannot be used interchangeably for body fat measurements.^{47,48}

Demographic and Socioeconomic Factors

Information on age, gender, monthly household income, and father's and mother's education was gathered via self-report.

Statistical Analysis

A series of intention-to-treat, 2-way repeated measures analysis of covariance were performed with intervention (ie, MI only, MI + PI, control) as the between-subject variable, time (pre-post) as the within-subjects variables and age, gender, and education of the father to examine group differences in change in physical activity, diet patterns and weight over the 12-month follow-up period. Paired *t* tests with Bonferroni adjustment were adopted for the post hoc analysis. Between-group effect size was calculated by using partial η squared. A *P* value of $<.05$ was considered statistically significant.

RESULTS

Descriptive statistics are shown in Table 1. Of the original 119 adolescents enrolled in each group, 113, 118 and 115 participants for the MI, MI + PI, and control groups, respectively, completed the 12-month assessment. Dropouts were not significantly different from completers on any baseline demographic or clinical variable.

A total of 357 obese adolescents participated in the study. The mean age in each group was 15.59, 15.57, and 15.78 years for the MI, MI + PI, and control groups, respectively. There were no significant differences between the 3 groups in terms of sociodemographic characteristics and anthropometric measurements at baseline.

We hypothesized (hypothesis bloc 1) that MI + PI would be effective in changing relevant weight loss outcomes compared with a passive control condition. Significant post hoc comparisons (Tables 2 and 3) were made for most of the comparisons. The MI + PI group was not superior in changing rates compared with the control condition for servings of vegetables (*P* = .65) and milk products (*P* = .27) per day, waist circumference (*P* = .51), and social functioning (*P* = .12). For all other psychometric variables (eg, school

TABLE 1 Sociodemographic Characteristics of the Obese Iranian Adolescents According to Study Group

Variable	MI (n = 119)	MI + PI (n = 119)	Control (n = 119)	P
Age, y	15.59 ± 1.31	15.57 ± 1.38	15.78 ± 1.19	.38
Gender, n (%)				
Male	55 (46.2)	53 (44.5)	65 (54.6)	.24
Female	64 (53.8)	66 (35.9)	54 (45.4)	
BMI, kg/m ²	33.07 ± 8.87	33.09 ± 5.86	32.92 ± 7.79	.95
BMI z score	2.83 ± 0.79	2.82 ± 0.62	2.75 ± 0.67	.83
BMI percentile	95.19 ± 4.72	94.63 ± 5.01	95.56 ± 4.63	.72
Mother's BMI, kg/m ²	39.46 ± 9.81	39.48 ± 8.87	39.31 ± 8.64	.77
Father's BMI, kg/m ²	36.22 ± 9.89	36.32 ± 9.83	36.02 ± 8.78	.83
Waist circumference, cm	102.59 ± 9.27	103.48 ± 8.19	95.51 ± 8.75	.94
Mother education, y	6.26 ± 3.42	6.19 ± 3.54	5.92 ± 3.89	.47
Father education, y	7.43 ± 3.86	7.28 ± 3.91	8.26 ± 4.21	.12
Household income (1000 rials) ^a	7434.63 ± 1891.39	7304.10 ± 2136.48	7364.67 ± 1944.10	.79
DXA, % fat	47.73 ± 7.42	47.70 ± 6.35	45.37 ± 8.31	.51
BIA	46.04 ± 4.03	45.85 ± 4.33	45.15 ± 4.72	.93

^a About 3200 rials = US \$1.00.

TABLE 2 Dietary and Anthropometric Outcomes According to Intervention Group for Baseline and Follow-up

Variable	Baseline				12-mo Follow-up				Time Effects				Post Hoc Test ^a	
	MI (n = 119)		MI + PI (n = 119)		MI (n = 113)		MI + PI (n = 118)		Time Effect		Time × Group Effect		MI + PI Versus Control MI	
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	F	P	η_p^2	F	P	P
Diet/food intake														
Total calories, kcal	3022.22 ± 1442.70	3000.52 ± 1449.14	2930.36 ± 1342.81	2785.21 ± 1240.31	2256.08 ± 1351.46	2919.95 ± 1471.92	58.09	<.001	0.14	25.03	<.001	0.12	.002	.032
Total dietary fat, g	105.55 ± 63.50	102.80 ± 60.85	95.23 ± 40.16	98.42 ± 49.83	74.40 ± 42.39	95.73 ± 40.10	28.74	<.001	0.75	15.77	<.001	0.09	.012	.041
Saturated fat, g	36.52 ± 23.70	36.29 ± 24.57	36.49 ± 23.75	32.54 ± 22.86	35.52 ± 21.48	30.03 ± 20.84	300.43	<.001	0.46	50.60	<.001	0.22	<.001	.029
Fruits and juice, servings/ d	1.27 ± 0.98	1.27 ± 0.97	1.21 ± 0.97	1.31 ± 0.96	1.33 ± 0.93	1.23 ± 0.97	2.85	.09	0.01	0.26	.77	0.00	.030	.17
Vegetables, servings/d	1.64 ± 0.76	1.66 ± 0.75	1.66 ± 0.73	1.71 ± 0.80	1.77 ± 0.76	1.64 ± 0.80	4.61	.03	0.01	2.24	.10	0.01	.65	.77
Snacks/desserts, servings/d	4.43 ± 1.52	4.33 ± 1.82	4.49 ± 1.63	4.12 ± 1.43	3.89 ± 1.65	4.55 ± 1.71	41.28	<.001	0.10	17.65	<.001	0.09	<.001	.04
Nondiet soda, servings/d	0.87 ± 0.66	0.85 ± 0.53	0.89 ± 0.42	0.77 ± 0.38	0.48 ± 0.35	0.95 ± 0.33	33.23	<.001	0.09	5.46	<.001	0.14	.001	.01
Milk, servings/ d	0.94 ± 0.49	0.95 ± 0.38	0.98 ± 0.48	0.95 ± 0.29	0.96 ± 0.43	0.97 ± 0.37	0.02	.89	0.00	0.08	.92	0.00	.27	.34
Fried foods, servings/d	0.81 ± 0.38	0.82 ± 0.33	0.84 ± 0.45	0.82 ± 0.30	0.63 ± 0.33	0.90 ± 0.44	6.45	.10	0.01	14.76	<.001	0.08	.01	.042
Anthropometry and body composition														
Total cholesterol, mmol/L	4.42 ± 0.73	4.40 ± 0.82	4.18 ± 0.96	4.41 ± 0.72	4.27 ± 0.78	4.20 ± 0.97	4.26	.04	0.01	4.70	.010	0.03	.021	.049
Triglycerides, mmol/L	1.01 ± 0.78	1.02 ± 0.77	1.05 ± 0.68	1.01 ± 0.77	0.86 ± 0.78	1.09 ± 0.72	5.68	.018	0.02	10.55	.001	0.06	.001	.031
BMI, kg/m ²	33.07 ± 8.87	33.09 ± 5.86	32.92 ± 7.79	32.44 ± 7.45	31.04 ± 6.46	32.95 ± 8.78	25.51	<.001	0.07	14.76	.001	0.08	.01	.032
BMI z score	2.83 ± 0.79	2.82 ± 0.62	2.75 ± 0.67	2.81 ± 0.76	2.58 ± 0.61	2.76 ± 0.70	12.73	<.001	0.04	12.13	<.001	0.06	.02	.048
DXA, % fat	47.73 ± 7.42	47.70 ± 6.35	45.37 ± 8.31	46.61 ± 7.91	45.96 ± 6.34	45.40 ± 7.45	44.49	<.001	0.11	13.47	<.001	0.07	.012	.38
BIA	46.04 ± 4.03	45.85 ± 4.33	45.15 ± 4.72	45.62 ± 4.32	42.32 ± 3.56	45.33 ± 4.86	13.45	<.001	0.04	11.77	<.001	0.06	.001	.38
Waist circumference, cm	102.59 ± 9.27	103.48 ± 8.19	95.51 ± 8.75	101.97 ± 9.11	101.70 ± 9.21	99.90 ± 8.26	2.67	.10	0.00	2.36	.09	0.01	.512	.717

Unless otherwise indicated, data are presented as mean ± SD. η_p^2 = partial η squared (measure of effect size).

^a Bonferroni adjustment.

functioning [$P = .03$], physical [$P = .01$] and psychosocial [$P = .01$] health, exercise [$P = .003$], dietary self-efficacy [$P < .001$]), dietary factors (eg, fat intake [$P = .01$], servings of snacks [$P < .001$] and fried food [$P = .01$], total calories [$P = .002$]), and anthropometric measures (eg, physical activity [$P = .001$], energy expenditure [$P = .001$], BMI [$P = .001$], cholesterol [$P = .02$], triglycerides [$P = .02$], DXA [$P = .01$], BIA [$P = .001$]), the comparison of MI + PI versus the control group yielded significant differences in favor of the intervention.

We further hypothesized (hypothesis bloc 2) that MI + PI would be superior in changing weight loss outcomes compared with MI only. The pattern of results was similar, albeit slightly weaker differences were found. For many outcome variables, significant post hoc differences emerged (Tables 2 and 3). The MI + PI group was not significantly superior compared with MI for adolescents only for servings of fruits ($P = .17$), vegetables ($P = .77$), and milk products ($P = .34$) per day; waist circumference ($P = .71$); DXA ($P = .38$); BIA ($P = .38$); and social ($P = .10$) and psychosocial ($P = .35$) functioning. For the other psychometric variables (eg, school functioning [$P = .02$], physical [$P = .04$], exercise [$P = .003$], dietary self-efficacy [$P < .001$]), dietary habits (eg, fat intake [$P = .04$], servings of snacks [$P = .04$] and fried food [$P = .04$], total calories [$P = .03$]), and anthropometric measures (eg, physical activity [$P = .01$], energy expenditure [$P = .003$], BMI [$P = .048$], cholesterol [$P = .049$], triglycerides [$P = .031$]), the comparison of MI + PI with MI only for adolescents exhibited significant differences.

DISCUSSION

To evaluate the role of parental involvement in MI weight loss interventions for obese adolescents, a prospective RCT was performed

TABLE 3 Energy Expenditure and Quality of Life Outcomes According to Intervention Group for Baseline and Follow-up

Variable	Baseline				12-mo Follow-up				Time Effects				Post Hoc Test ^a				
	MI (n = 119)		MI + PI (n = 119)		Control (n = 119)		MI (n = 113)		MI + PI (n = 118)		Control (n = 115)		Time Effect	Time × Group Effect	MI + PI Versus Control	MI + PI Versus MI	
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	F	P	η _p ²	F					P
Measure																	
CDSS	6.16 ± 2.46	6.14 ± 2.51	6.02 ± 2.32	6.88 ± 2.72	7.11 ± 2.62	5.71 ± 2.82	6.88 ± 2.72	7.11 ± 2.62	5.71 ± 2.82	34.56	<.001	0.09	23.59	<.001	0.12	<.001	.01
WEL	87.10 ± 27.37	87.29 ± 28.94	93.61 ± 29.14	101.27 ± 27.23	108.05 ± 24.28	93.21 ± 29.60	101.27 ± 27.23	108.05 ± 24.28	93.21 ± 29.60	45.01	<.001	0.11	12.51	<.001	0.07	.002	.04
PES	5.61 ± 1.74	5.59 ± 1.87	5.69 ± 1.82	6.75 ± 1.15	6.68 ± 1.97	5.39 ± 2.18	6.75 ± 1.15	6.68 ± 1.97	5.39 ± 2.18	70.12	<.001	0.16	38.01	<.001	0.17	.003	.01
Self-reported PA length	0.84 ± 0.49	0.86 ± 0.45	0.70 ± 0.35	1.12 ± 0.61	1.30 ± 0.66	0.66 ± 0.32	1.12 ± 0.61	1.30 ± 0.66	0.66 ± 0.32	35.16	<.001	0.09	13.43	<.001	0.07	.004	.01
Self-reported energy expenditure	247.73 ± 160.36	220.50 ± 106.63	271.87 ± 127.99	434.07 ± 128.98	509.07 ± 128.15	268.14 ± 109.08	434.07 ± 128.98	509.07 ± 128.15	268.14 ± 109.08	115.08	<.001	0.54	31.92	<.001	0.39	.002	.02
Objective PA length	0.67 ± 0.38	0.68 ± 0.28	0.65 ± 0.39	0.98 ± 0.37	0.59 ± 0.40	1.34 ± 0.42	0.98 ± 0.37	0.59 ± 0.40	1.34 ± 0.42	36.26	<.001	0.09	17.06	<.001	0.09	.001	.01
Objective energy expenditure	280.17 ± 82.83	274.90 ± 85.32	293.95 ± 81.83	500.02 ± 94.94	680.08 ± 88.14	262.92 ± 99.22	500.02 ± 94.94	680.08 ± 88.14	262.92 ± 99.22	82.65	<.001	0.59	33.67	<.001	0.54	.001	.003
Measure																	
Total score	63.19 ± 20.58	62.46 ± 20.18	64.74 ± 19.91	64.34 ± 21.11	65.14 ± 18.88	63.73 ± 17.16	64.34 ± 21.11	65.14 ± 18.88	63.73 ± 17.16	3.05	.08	0.01	4.081	.01	0.02	.21	.31
Physical health	66.71 ± 19.35	66.17 ± 19.28	65.12 ± 24.53	70.13 ± 19.20	71.84 ± 20.88	62.93 ± 25.85	70.13 ± 19.20	71.84 ± 20.88	62.93 ± 25.85	8.55	<.001	0.02	8.97	<.001	0.05	.01	.04
Psychosocial health	67.69 ± 17.19	67.04 ± 18.49	67.54 ± 19.3	69.08 ± 21.62	70.16 ± 15.09	66.30 ± 16.83	69.08 ± 21.62	70.16 ± 15.09	66.30 ± 16.83	4.26	.04	0.01	5.85	<.001	0.04	.01	.35
Emotional functioning	68.97 ± 19.34	68.35 ± 20.42	69.59 ± 19.92	71.64 ± 19.99	71.81 ± 21.79	68.27 ± 21.82	71.64 ± 19.99	71.81 ± 21.79	68.27 ± 21.82	7.15	.008	0.02	6.20	.002	0.03	.03	.24
Social functioning	69.84 ± 18.41	69.12 ± 19.74	70.12 ± 17.42	70.84 ± 19.08	69.12 ± 19.47	70.01 ± 17.98	70.84 ± 19.08	69.12 ± 19.47	70.01 ± 17.98	3.49	.06	0.01	3.48	.03	0.02	.12	.10
School functioning	65.01 ± 15.79	64.18 ± 16.78	68.83 ± 20.63	66.80 ± 17.62	71.26 ± 18.20	65.73 ± 23.77	66.80 ± 17.62	71.26 ± 18.20	65.73 ± 23.77	5.40	.02	0.01	12.70	<.001	0.07	.03	.02

^a η_p^2 = partial η squared (measure of effect size); PA = physical activity.

^a Bonferroni adjustment.

comparing 3 intervention groups: MI for adolescents only, MI for adolescents with parental involvement, and a passive control group. Anthropometric, biochemical, psychosocial, and behavioral data were collected at baseline and 12 months later. Results revealed significant effects for MI with parental involvement compared with the control group (hypothesis bloc 1) and compared with MI for adolescents only (hypothesis bloc 2). This pattern could be shown for most of the anthropometric, biochemical, psychosocial, and behavioral outcome variables supporting the hypotheses.

The present results are in line with findings of a recent meta-analysis regarding MI interventions for different types of health behaviors, as well as interventions targeting the obese adolescent population directly.²³ The additional effectiveness of parental involvement support the associations found in the meta-analysis. According to Gayes et al,²³ studies that assessed MI with both adolescent and parental input reported the largest effects, demonstrating that conducting MI with both parent and adolescent is more effective than conducting MI with either parent or adolescent only. Previous behavioral interventions for obese adolescents were inconclusive regarding the role of parental involvement.^{24–26} The present study can strengthen the evidence for parental involvement in the context of an MI intervention.

Some study limitations must be acknowledged. Despite the appropriate sample size and the low dropout rate in an understudied population, the results might be limited to Persian culture. Thus, generalizability to other study contexts should be made with caution, although the present findings are in line with evidence obtained in Western samples.²¹ However, more studies are needed to directly

compare the effects of parental involvement and MI among samples from different cultures. In addition, we did not use objective outcome physical activity measures for the whole sample. Furthermore, intervention was delivered by trained interventionists; consecutive studies are needed to validate the replicability of our findings with other interventionists but also with other nonspecialist interventionists such as trained teachers (coached according to the MI training manual). Finally, the present study was conducted by using a 12-month follow-up. A subsequent next step would be to investigate the longer term effects of the intervention (ie, after several years).

CONCLUSIONS

Because our study was a clinical assessment, future studies should replicate our findings in real-world settings, such as integrating MI as a brief strategy into the daily routines of a school, particularly for obese adolescents, but for nonobese adolescents as a prevention tool as well. Effects of parental involvement should be tested via invitation letters sent by the school and brief training sessions, in which adolescents and their parents could be motivated to change their dietary and physical activity habits.

Future studies might extend the investigation to different types of the delivery mode, not only involving a parent but also varying the intensity level of parental involvement because this dose-response relation has been examined by previous meta-analyses. Beyond that, examination of subgroup effects would provide further insights into the topic; for example, the effects of parental involvement might vary according to the gender, age, quality of the relationship between adolescent and parent, or the family's cooking habits.⁴⁹

The present study evaluated and compared the role of parental

involvement in MI interventions for obese adolescents compared with MI for adolescents only and control subjects. The results show that MI with parental involvement is an effective strategy in changing obesity-related outcomes and has additional effects beyond MI with adolescents only, which is important when administering MI interventions in school settings.

REFERENCES

- Schmidt Morgen C, Rokholm B, Sjöberg Brixval C, et al. Trends in prevalence of overweight and obesity in Danish infants, children and adolescents—are we still on a plateau? *PLoS One*. 2013;8(7):e69860
- Moraes AC, Falcão MC. Lifestyle factors and socioeconomic variables associated with abdominal obesity in Brazilian adolescents. *Ann Hum Biol*. 2013;40(1):1–8
- Lobstein T, Baur L, Uauy R; IASO International Obesity Task Force. Obesity in children and young people: a crisis in public health. *Obes Rev*. 2004;5(suppl 1): 4–104
- World Health Organization. *The Challenge of Obesity—Quick Statistics*. 2012 ed. Geneva, Switzerland: World Health Organization; 2012. Available at: www.euro.who.int/en/health-topics/noncommunicable-diseases/obesity/data-and-statistics. Accessed January 5, 2015
- Raj M, Kumar RK. Obesity in children & adolescents. *Indian J Med Res*. 2010;132: 598–607
- Moayeri H, Bidad K, Aghamohammadi A, et al. Overweight and obesity and their associated factors in adolescents in Tehran, Iran, 2004-2005. *Eur J Pediatr*. 2006;165(7):489–493
- Behzadnia S, Vahidshahi K, Hamzeh Hosseini S, Anvari S, Ehteshami S. Obesity and related factors in 7-12 year-old elementary school students during 2009-2010 in Sari, Iran. *Med Glas (Zenica)*. 2012;9(1):86–90
- Centers for Diseases Control and Prevention. *Childhood Obesity Facts*. Atlanta, GA: Centers for Diseases Control and Prevention; 2012
- Puhl R, Brownell KD. Bias, discrimination, and obesity. *Obes Res*. 2001;9(12): 788–805
- Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002; 360(9331):473–482
- Budd GM, Hayman LL, Crump E, et al. Weight loss in obese African American and Caucasian adolescents: secondary analysis of a randomized clinical trial of behavioral therapy plus sibutramine. *J Cardiovasc Nurs*. 2007;22(4):288–296
- Pimenta F, Leal I, Maroco J, Ramos C. Brief cognitive-behavioral therapy for weight loss in midlife women: a controlled study with follow-up. *Int J Womens Health*. 2012;4:559–567
- Jelalian E, Boergers J, Alday CS, Frank R. Survey of physician attitudes and practices related to pediatric obesity. *Clin Pediatr (Phila)*. 2003;42(3):235–245
- Befort CA, Nollen N, Ellerbeck EF, Sullivan DK, Thomas JL, Ahluwalia JS. Motivational interviewing fails to improve outcomes of a behavioral weight loss program for obese African American women: a pilot randomized trial. *J Behav Med*. 2008;31(5):367–377
- Hardcastle SJ, Taylor AH, Bailey MP, Harley RA, Hagger MS. Effectiveness of a motivational interviewing intervention on weight loss, physical activity and cardiovascular disease risk factors: a randomised controlled trial with a 12-month post-intervention follow-up. *Int J Behav Nutr Phys Act*. 2013;10:40
- Wong EM, Cheng MM. Effects of motivational interviewing to promote weight loss in obese children. *J Clin Nurs*. 2013;22(17–18):2519–2530
- Martins RK, McNeil DW. Review of motivational interviewing in promoting health behaviors. *Clin Psychol Rev*. 2009; 29(4):283–293
- Miller WR, Rollnick S. *Motivational Interviewing: Preparing People for Change*. 2nd ed. New York, NY: Guilford Press; 2002
- Levensky ER, Forcehimes A, O'Donohue WT, Beitz K. Motivational interviewing: an evidence-based approach to counseling helps patients follow treatment recommendations. *Am J Nurs*. 2007; 107(10):50–58, quiz 58–59
- Rollnick S, Butler CC, McCambridge J, Kinnersley P, Elwyn G, Resnicow K. Consultations about changing behaviour. *BMJ*. 2005;331(7522):961–963

21. Sormunen M, Tossavainen K, Turunen H. Finnish parental involvement ethos, health support, health education knowledge and participation: results from a 2-year school health intervention. *Health Educ Res.* 2013;28(2):179–191
22. Davidson KW, Goldstein M, Kaplan RM, et al. Evidence-based behavioral medicine: what is it and how do we achieve it? *Ann Behav Med.* 2003;26(3): 161–171
23. Gayes LA, Steele RG. A meta-analysis of motivational interviewing interventions for pediatric health behavior change. *J Consult Clin Psychol.* 2014;82(3):521–535
24. van der Kruk JJ, Kortekaas F, Lucas C, Jager-Wittenaar H. Obesity: a systematic review on parental involvement in long-term European childhood weight control interventions with a nutritional focus. *Obes Rev.* 2013;14(9):745–760
25. Van Lippevelde W, Verloigne M, De Bourdeaudhuij I, et al. Does parental involvement make a difference in school-based nutrition and physical activity interventions? A systematic review of randomized controlled trials. *Int J Public Health.* 2012;57(4):673–678
26. Hingle MD, O'Connor TM, Dave JM, Baranowski T. Parental involvement in interventions to improve child dietary intake: a systematic review. *Prev Med.* 2010;51(2):103–111
27. Gourlan M, Sarrazin P, Trouilloud D. Motivational interviewing as a way to promote physical activity in obese adolescents: a randomised-controlled trial using self-determination theory as an explanatory framework. *Psychol Health.* 2013;28(11):1265–1286
28. Lloyd-Richardson EE, Jelalian E, Sato AF, Hart CN, Mehlenbeck R, Wing RR. Two-year follow-up of an adolescent behavioral weight control intervention. *Pediatrics.* 2012;130(2). Available at: www.pediatrics.org/cgi/content/full/130/2/e281
29. Walpole B, Dettmer E, Morrongiello BA, McCrindle BW, Hamilton J. Motivational interviewing to enhance self-efficacy and promote weight loss in overweight and obese adolescents: a randomized controlled trial. *J Pediatr Psychol.* 2013; 38(9):944–953
30. Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, et al. 2000 CDC growth charts for the United States: methods and development. *Vital Health Stat.* 2002 (246):1–190
31. Burke BL, Arkowitz H, Menchola M. The efficacy of motivational interviewing: a meta-analysis of controlled clinical trials. *J Consult Clin Psychol.* 2003;71(5): 843–861
32. National Health and Medical Research Council. *Food for Health: Dietary Guidelines for Children and Adolescents in Australia.* Canberra, Australia: Commonwealth Department of Health and Ageing; 2003
33. World Health Organization. *Global Recommendations on Physical Activity for Health.* Geneva, Switzerland: World Health Organization; 2010
34. Moyers TB, Martin T, Manuel JK, Miller WR, Ernst D. *Motivational Interviewing Treatment Integrity 3.1.1: Revised Global Scales.* Albuquerque, NM: University of New Mexico; 2010
35. Parcel GS, Edmundson E, Perry CL, et al. Measurement of self-efficacy for diet-related behaviors among elementary school children. *J Sch Health.* 1995;65(1): 23–27
36. Clark MM, Abrams DB, Niaura RS, Eaton CA, Rossi JS. Self-efficacy in weight management. *J Consult Clin Psychol.* 1991;59(5):739–744
37. Navidian A, Abedi M, Baghban I, Fatehizadeh M. Reliability and validity of the Weight Efficacy Lifestyle questionnaire in overweight and obese individuals. *JBS.* 2009;3(3):217–222
38. Rockett HR, Breitenbach M, Frazier AL, et al. Validation of a youth/adolescent food frequency questionnaire. *Prev Med.* 1997;26(6):808–816
39. Gross LD, Sallis JF, Buono MJ, Roby JJ, Nelson JA. Reliability of interviewers using the seven-day physical activity recall. *Res Q Exerc Sport.* 1990;61(4):321–325
40. Vashghani-Farahani A, Tahmasbi M, Asheri H, Ashraf H, Nedjat S, Kordi R. The Persian, last 7-day, long form of the International Physical Activity Questionnaire: translation and validation study. *Asian J Sports Med.* 2011;2(2): 106–116
41. Crouter SE, Horton M, Bassett DR Jr. Use of a two-regression model for estimating energy expenditure in children. *Med Sci Sports Exerc.* 2012;44(6):1177–1185
42. Pakpour AH. Psychometric properties of the Iranian version of the Pediatric Quality of Life Inventory™ Short Form 15 Generic Core Scales. *Singapore Med J.* 2013;54(6):309–314
43. Chen H-J, Wang Y. Use of percentiles and z -scores in anthropometry. In: Preedy V, ed. *Handbook of Anthropometry: Physical Measures of Human Form in Health and Disease.* London, UK: Springer; 2012:29–48
44. Houtkooper LB, Lohman TG, Going SB, Howell WH. Why bioelectrical impedance analysis should be used for estimating adiposity. *Am J Clin Nutr.* 1996;64(suppl 3):436S–448S
45. Malavolti M, Mussi C, Poli M, et al. Cross-calibration of eight-polar bioelectrical impedance analysis versus dual-energy x-ray absorptiometry for the assessment of total and appendicular body composition in healthy subjects aged 21–82 years. *Ann Hum Biol.* 2003;30(4): 380–391
46. Demura S, Sato S, Kitabayashi T. Percentage of total body fat as estimated by three automatic bioelectrical impedance analyzers. *J Physiol Anthropol Appl Human Sci.* 2004;23(3): 93–99
47. Eisenmann JC, Heelan KA, Welk GJ. Assessing body composition among 3- to 8-year-old children: anthropometry, BIA, and DXA. *Obes Res.* 2004;12(10): 1633–1640
48. Ellis KJ. Measuring body fatness in children and young adults: comparison of bioelectric impedance analysis, total body electrical conductivity, and dual-energy x-ray absorptiometry. *Int J Obes Relat Metab Disord.* 1996;20(9):866–873
49. Bauer KW, Neumark-Sztainer D, Hannan PJ, Fulkerson JA, Story M. Relationships between the family environment and school-based obesity prevention efforts: can school programs help adolescents who are most in need? *Health Educ Res.* 2011;26(4):675–688

Motivational Interviewing With Parents for Obesity: An RCT
Amir H. Pakpour, Paul Gellert, Stephan U. Dombrowski and Bengt Fridlund
Pediatrics 2015;135:e644; originally published online February 9, 2015;
DOI: 10.1542/peds.2014-1987

Updated Information & Services	including high resolution figures, can be found at: /content/135/3/e644.full.html
Supplementary Material	Supplementary material can be found at: /content/suppl/2015/02/04/peds.2014-1987.DCSupplemental.html
References	This article cites 40 articles, 3 of which can be accessed free at: /content/135/3/e644.full.html#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Psychiatry/Psychology /cgi/collection/psychiatry_psychology_sub Obesity /cgi/collection/obesity_new_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: /site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: /site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Motivational Interviewing With Parents for Obesity: An RCT

Amir H. Pakpour, Paul Gellert, Stephan U. Dombrowski and Bengt Fridlund

Pediatrics 2015;135:e644; originally published online February 9, 2015;

DOI: 10.1542/peds.2014-1987

The online version of this article, along with updated information and services, is located on the World Wide Web at:

</content/135/3/e644.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

